

(CLAW, \vec{C}_3)-FREE DIGRAPHS WITH UNBOUNDED DICHROMATIC NUMBER

GUILLAUME AUBIAN AND LUIS KUFFNER

ABSTRACT. We construct orientations of rook graphs (whose underlying graphs are claw-free) that contain no \vec{C}_3 but have unbounded dichromatic number. This disproves a conjecture of Aboulker, Charbit and Naserasr and improves a result of Carbonero, Koerts, Moore and Spirkel.

1. INTRODUCTION AND NOTATION

A hero is a tournament H such that tournaments not containing H have bounded dichromatic number. In a seminal paper [3], Berger, Choromanski, Chudnovski, Fox, Loebl, Scott, Seymour and Thomassé obtained a structure theorem for heroes. Harutyunyan, Le, Newman and Thomassé [7] later proved that oriented graphs with bounded independence number (tournaments being the special case where the independence number equals one) not containing a hero have bounded dichromatic number. Chudnovsky, Scott and Seymour [6] subsequently showed that for any transitive tournament H and oriented star S , oriented graphs containing neither S nor H as induced subdigraphs have bounded dichromatic number. Aboulker, Charbit and Naserasr [2] thus wondered whether (and actually conjectured that), for any hero H and any oriented star forest S , oriented graphs containing neither S nor H as induced subdigraphs have bounded dichromatic number.

Aboulker, Aubian and Charbit [1] provided a counterexample when $S = K_1 + \vec{K}_2$. Carbonero, Koerts, Moore and Spirkel [5] did so when $H = \Delta(1, 1, C_3)$ and S is any orientation of $K_{1,4}$, and also when $H = \vec{C}_3$ and S is any orientation of $K_{1,5}$. We improve these two latter results with a counterexample when $H = \vec{C}_3$ and S is any orientation of $K_{1,3}$, i.e. the claw.

Definitions in this paper are standard and follow from Bondy and Murty's book [4]. In this paper, we study oriented graphs, i.e. directed graphs with no pair of opposite arcs. Given two integers $n, i \geq 0$, we write n_i for the i th least significant bit of the binary expansion of n , starting from $i = 0$.

2. CONSTRUCTION AND PROOF

Let N be a positive integer. The $N \times N$ *rook graph* R_N is the graph with vertex set $[1, N]^2$ and with an edge between two vertices whenever they share their first coordinate or they share their second coordinate. Intuitively, it is the graph describing how a rook moves on a $N \times N$ chessboard. It is the line graph of $K_{N,N}$, and therefore is claw-free. We orient it into a digraph D_N as follows. For any vertices (a, b) , (a, d) , (c, b) , we orient the edge from (a, b) to (a, d) if and only if $b_i = a_i$, where $i = \min\{j \mid b_j \neq d_j\}$, and we orient the edge from (a, b) to (c, b) if and only if $b_i \neq a_i$, where $i = \min\{j \mid a_j \neq c_j\}$.

Proposition 1. *The digraph D_N contains no \vec{C}_3 .*

Proof. Given three vertices (a, b) , (a, c) and (a, d) on the same row, let $i = \min\{j \mid \{b_j, c_j, d_j\} = \{0, 1\}\}$. Without loss of generality let us assume that $b_i \neq c_i = d_i$. Edges $(a, b)(a, c)$ and $(a, b)(a, d)$ are oriented only depending on whether $a_i = b_i$ and thus have the same orientation. Thus, (a, b) , (a, c) and (a, d) can not form a \vec{C}_3 . The same argument applies to any column. Since every triangle in a rook graph is contained entirely in a single row or in a single column, D_N contains no \vec{C}_3 . \square

Lemma 1. *When $|a - c| = |b - d|$, vertices (a, b) , (a, d) , (c, d) , (c, b) induce a directed 4-cycle.*

Date: November 2025.

Proof. Let $i = \min\{j \mid |a - c|_j = 1\}$. Then $i = \min\{j \mid a_j \neq c_j\} = \min\{j \mid b_j \neq d_j\}$. Up to permuting a and c and/or permuting b and d , we may assume $a_i = b_i = 1$ and $c_i = d_i = 0$. Since $b_i = a_i$, we have $(a, b) \rightarrow (a, d)$. Since $d_i \neq a_i$, we have $(a, d) \rightarrow (c, d)$. Since $d_i = c_i$, we have $(c, d) \rightarrow (c, b)$. Since $b_i \neq c_i$, we have $(c, b) \rightarrow (a, b)$. Thus the four vertices form the directed cycle $(a, b) \rightarrow (a, d) \rightarrow (c, d) \rightarrow (c, b) \rightarrow (a, b)$. \square

Note that it actually suffices that $d_2(a, c) = d_2(b, d)$ where d_2 is the dyadic distance, but this stronger condition suffices for the sake of our result.

Theorem 1. *The family $(D_N)_{N \geq 1}$ has unbounded dichromatic number.*

Proof. Fix k . The Gallai–Witt theorem [8] (also known as the multidimensional van der Waerden theorem) implies that there exists N such that any k -colouring of $[1, N]^2$ contains a monochromatic axis-parallel square, that is, four points of the form (x, y) , $(x + t, y)$, $(x, y + t)$, $(x + t, y + t)$ for some integers $x, y, t > 0$. Thus, for N large enough, any colouring of $V(D_N)$ with k colours contains a monochromatic square as above. By the previous lemma, the four vertices of such a square induce a directed 4-cycle. Hence, some colour class does not induce an acyclic subdigraph. Since k is arbitrary, the dichromatic number of D_N is unbounded. \square

REFERENCES

1. Pierre Aboulker, Guillaume Aubian, and Pierre Charbit, *Heroes in oriented complete multipartite graphs*, Journal of Graph Theory **105** (2024), no. 4, 652–669.
2. Pierre Aboulker, Pierre Charbit, and Reza Naserasr, *Extension of gyárfás–sumner conjecture to digraphs*, The Electronic Journal of Combinatorics **28** (2021), no. 2, P2.27.
3. Eli Berger, Krzysztof Choromanski, Maria Chudnovsky, Jacob Fox, Martin Loebl, Alex Scott, Paul Seymour, and Stéphan Thomassé, *Tournaments and colouring*, Journal of Combinatorial Theory, Series B **103** (2013), no. 1, 1–20.
4. J.A. Bondy and U.S.R Murty, *Graph theory*, 1st ed., Springer Publishing Company, Incorporated, 2008.
5. Alvaro Carbonero, Hidde Koerts, Benjamin Moore, and Sophie Spirkl, *On heroes in digraphs with forbidden induced forests*, European Journal of Combinatorics **125** (2025), 104104.
6. Maria Chudnovsky, Alex Scott, and Paul Seymour, *Induced subgraphs of graphs with large chromatic number. xi. orientations*, European Journal of Combinatorics **76** (2019), 53–61.
7. Ararat Harutyunyan, Tien-Nam Le, Alantha Newman, and Stéphan Thomassé, *Coloring dense digraphs*, Electronic Notes in Discrete Mathematics **61** (2017), 577–583.
8. Ernst Witt, *Ein kombinatorischer satz der elementargeometrie*, Mathematische Nachrichten **6** (1951), 261–262.

CRED, UNIVERSITÉ PARIS-PANTHÉON-ASSAS

UNIVERSITÉ PARIS CITÉ, CNRS, IRIF, F-75013 PARIS, FRANCE